ISSN: 2301-5690

INTERNATIONAL CONFERENCE



The Second International Conference on Engineering and Technology Development

2ªICETD 2013

27, 28, 29 August 2013, Bandar Lampung, Indonesia

PROCEEDINGS







Bergelander alle interdiet die bestelie in die best Bestelie in die bestelie i



Hosted by : Faculty of Engineering and Faculty of Computer Science, Bandar Lampung University (UBL), Indonesia

ZndICETD 2013

THE SECOND INTERNATIONAL CONFERENCE ON ENGINEERING AND TECHNOLOGY DEVELOPMENT

> 28 -30 January 2013 Bandar Lampung University (UBL) Lampung, Indonesia

PROCEEDINGS

Organized by:



Faculty of Computer Science and Faculty of Engineering Bandar Lampung University (UBL) JI. Zainal Abidin Pagar Alam No.89 Labuhan Ratu, Bandar Lampung, Indonesia Phone: +62 721 36 666 25, Fax: +62 721 701 467 website :www.ubl.ac.id

PREFACE

The Activities of the International Conference is in line and very appropriate with the vision and mission of Bandar Lampung University (UBL) to promote training and education as well as research in these areas.

On behalf of the Second International Conference on Engineering and Technology Development (2^{nd} ICETD 2013) organizing committee, we are very pleased with the very good response especially from the keynote speaker and from the participans. It is noteworthy to point out that about 80 technical papers were received for this conference.

The participants of the conference come from many well known universities, among others : University Kebangsaan Malaysia - Malaysia, APTIKOM - Indonesia, Institut Teknologi sepuluh November - Indonesia, Surya Institute - Indonesia, International Islamic University - Malaysia, STMIK Mitra Lampung - lampung, Bandung Institut of Technology - Bandung, Lecture of The Malahayati University, B2TP - BPPT Researcher - lampung, Starch Technology Center - Lampung, Universitas Islam Indonesia – Indonesia, Politeknik Negeri Malang Malang, University of Kitakyushu – Japan, Gadjah Mada University – Indonesia, Universitas Malahayati – Lampung, Lampung University – lampung, Starch Technology Center - Lampung, Universitas Riau - Riau, Hasanuddin University -Indonesia, Diponegoro University – Indonesia, King Abdulaziz University – Saudi Arabia, Parahyangan Catholic University – Indonesia, National Taiwan University-Taiwan, Surakarta Christian University – Indonesia, Sugijapranata Catholic University - Indonesia, Semarang University - Indonesia, University of Brawijaya -Indonesia, PPKIA Tarakanita Rahmawati – Indonesia, Kyushu University, Fukuoka - Japan, Science and Technology Beijing - China, Institut Teknologi Sepuluh Nopember – Surabaya, Researcher of Starch Technology Center, Universitas Muhammadiyah Metro – Metro, National University of Malaysia – Malaysia.

I would like to express my deepest gratitude to the International Advisory Board members, sponsor and also to all keynote speakers and all participants. I am also gratefull to all organizing committee and all of the reviewers who contribute to the high standard of the conference. Also I would like to express my deepest gratitude to the Rector of Bandar Lampung University (UBL) who give us endless support to these activities, so that the conference can be administrated on time

Bandar Lampung, 29 August 2013-08-26

Mustofa Usman, Ph.D 2nd ICETD Chairman

PROCEEDINGS

2nd ICETD 2013

The Second International Conference On Engineering And Technology Development

28 - 30 January 2013

INTERNATIONAL ADVISORY BOARD

Y. M Barusman, Indonesia Ahmad F. Ismail, Malaysia Mustofa Usman, Indonesia Moses L. Singgih, Indonesia Andreas Dress, Germany Faiz A.M Elfaki, Malaysia Warsono, Indonesia Raihan Othman, Malaysia Zeng Bing Zen, China Tjin Swee Chuan, Singapore Khomsahrial R, Indonesia Rony Purba, Indonesia Alex Tribuana S, Indonesia Hon Wei Leong, Singapore Imad Khamis, USA Rozlan Alias, Malaysia Rudi Irawan. Indonesia Gusri Ibrahim, Indonesia Jamal I Daoud, Malaysia Riza Muhida, Indonesia Heri Riyanto, Indonesia Agus Wahyudi, Indonesia Lilies Widojoko, Indonesia

PROCEEDINGS

2nd ICETD 2013

The Second International Conference On Engineering And Technology Development

28 - 30 January 2013

STEERING COMMITTEE

Executive Advisors Dr. M. Yusuf S. Barusman Andala R. P. Barusman, MA.Ec

> **Chairman** Mustofa Usman, Ph.D

Co-Chairman Dr. Ir. Hery Riyanto, MT Ahmad Cucus, S.Kom., M.Kom

Secretary Marzuki, S.Kom., M.Kom Maria Shusanti Febrianti, S.Kom., M.Kom

Technical Committee

Indyah Kumoro, ST. IAI Ardiansyah, ST., MT Sofiah Islamiah, ST. MT Taqwan Thamrin, ST., MSc Dina Ika Wahyuningsih, S.Kom Agus Sukoco, M.Kom Hj. Susilowati, ST. MT Haris Murwadi, ST, MT Robby Yuli Endra, S.Kom., M.Kom Fenty Ariani, S.Kom., M.Kom

Treasure

Samsul Bahri, SE Dian Agustina, SE

PROCEEDINGS

2nd ICETD 2013

The Second International Conference On Engineering And Technology Development

28 - 30 January 2013

ORGANIZING COMMITTEE

Chair Person Dr. Ir. Hery Riyanto, MT

Vice Chair Person Yuthsi Aprilinda, S.Kom., M.Kom

> **Treasure** Dian Agustina, S.E

Secretary Aprizal, ST. MT Ir. Tjejeng Sofyan, MM Ir. Muhammad Zein, MT Ir. Bambang Pratowo, MT

Special Events

Ir. Juniardi, MT Ir. Indra Surya, MT Ir. Sugito, MT DR. Baginda Simaibang, M.Ed Berry Salatar, S.Pd Yanuar Dwi Prasetyo, S.Pd.I., M.A

Receiptionist

Ir. Najamudin, MT Kunarto, ST. MT IB. Ilham Malik, ST. MT Ir.A Ikhsan Karim, MT Ir. Asikin, MT Usman Rizal, ST., M.MSi

Transportation and Acomodation

Irawati, SE Desi Puspita Sari, S.E Tanto Lailam, S.H 2nd International Conference on Engineering and Technology Development (ICETD 2013) Universitas Bandar Lampung Faculty of Engineering and Faculty of Computer Science

Ilyas Sadad, S.T., M.T

Publication and Documentation

Ir. Indriati Agustina Gultom, M.M Noning Verawati, S.Sos Hesti, S.H Rifandi Ritonga, SH Violita, S.I.Kom

Cosumption

Dra. Yulfriwini, M.T Wiwin Susanty, S.Kom., M.Kom Fenty Ariani, S.Kom., M.Kom Reni Nursyanti, S.Kom., M.Kom Erlangga, S.Kom Arnes Yuli Vandika, S.Kom

Facility and Decoration

Siti Rahma Wati,SE Dina Ika Wahyuningsih, S.Kom Zainal Abidin, SE Ahyar Saleh, SE Eko Suhardiyanto Wagino Sugimin

Table Of Content

Drganizing Committee Γable Of Content	i v
Keynote Speaker	
Recent Advances in Biofuel Cell and Emerging Hybrid System Abdul Aziz Ahmad and Raihan Othman	1
2. Waste Utilization Study Tailing Gold Mine in Way Linggo-Lampung, as Fin Aggregate Materials for Producing Mortar Materials based on concept of Green Technology Lilies Widojoko & Susilawati.	e 1 8
 Infrastructure Health Monitoring System (SHM) Development, a Necessity fo Maintance and Investigation Prof. Dr. Priyo Suprobo, Faimun, Arie Febry	r 7
4. Four Phases Quality Function Deployment (Qfd) By Considering Kano Concept Time And Manufacturing Cost Prof. Dr. Moses L Singgih, Dyah L. Trenggonowati, Putu D. Karningsih 2	, 2

Speaker

1.	Comparative Analysis for The Multi Period Degree Minimum Spanning Tree Problem
	Wamiliana, Amanto, and Mustofa Usman
2.	Choosing The Right Software In Supporting The Successful of Enterprise ERP Implementation Yodhie Yuniarthe, Idris Asmuni
3.	Climate Adaptive Technology In Maintaining Vernacularism Of Urban Kampong Case study: KampungAdat (Indiginous) Mahmud, Bandung District,West Java Marcus Gartiwa
4.	The Prospect Of Diesohol In Facing Fossil Fuel Crissis M.C. Tri Atmodjo
5.	The Potential Of Agriculture And Forestry Biomass Wastes As Source Of Bioenergy Hardoyo
6.	The Importance of Education Facility as Sustainable Urban Generation Tool Fritz Akhmad Nuzir, Haris Murwadi and Bart Julien Dewancker
7.	The implementation of Secton Method for Solving Systems of Non Linear Equations Nur Rokhman
8.	Quality Control Analysis Into Decrease The Level Defects On Coffee Product Heri Wibowo, Sulastri and Emy Khikmawati
9.	Public Transportion Crisis In Bandar Lampung Ida Bagus Ilham Malik
10	 Geospatial Analysis of Land Use Change in Way Kuripan Watershed, Bandar Lampung City Candra Hakim Van Rafi'i1., Dyah Indriana Kusumastuti2., Dwi Jokowinarno
11	. Material Utilization Technology Of Agriculture And Forestry Waste Hardoyo
12	. The Supply Chain System Of Cassava On The Tapioca Industry Hardoyo
13	. Glass Technology In Natural Light Glasses On Aperture Element In The Architecture World Muhammad Rija & MT Pedia Aldy

14. An Eksperimental Permeable Asphalt Pavement Using Local Material Domato Stone On Quality Of Porous Asphalt		
Firdaus Chairuddin, Wihardi Tjaronge, Muhammad Ramli, Johannes Patanduk		
 Coordination Of Architectural Concepts And Construction Systems Eddy Hermanto. 129 		
 Seismic Assessment of RC Building Using Pushover Analysis Riza Ainul Hakim. 136 		
 Viscosity and Liquidity Index Relation for Elucidating Mudflow Behavior Budijanto Widjaja and Shannon Hsien-Heng Lee. 		
18. The Use of Pozzolanic Material for Improving Quality of Strontium Liquid Waste Cementation in Saline Environment during Nuclear Waste Immobilization Process		
Muhammad Yusuf, HayuTyasUtami, Tri SulistiyoHariNugroho, SusetyoHarioPutero		
 Geospatial Analysis Of Land Use And Land Cover Changes For Discharge At Way Kualagaruntang Watershed In Bandar Lampung Fieni Yuniarti, Dyah Indriana K, Dwi Joko Winarno		
20. Wifi Network Design For High Performance Heru Nurwarsito, , KasyfulAmron, BektiWidyaningsih		
 Studi on The Efficiency Using Nature Materials in The Structural Elements of Reinforced Concrete Beam Yasser, Herman Parung, M. Wihardi Tjaronge, Rudy Djamaluddin 167 		
 The Research Of Slow Release Nitrogen Fertilizer Applied In Sugarcane (Saccharum Officinarum) For Green Energy Bioethanol M.C. Tri Atmodjo, Agus Eko T. Nurul Rusdi, Sigit Setiadi, and Rina 179 		
23. Energy Utilization Technology Of Agriculture And Forestry Waste Hardoyo		
 Implementation Of Fuzzy Inference System With Tsukamoto Method For Study Programme Selection Fenty Ariani and Robby Yuli Endra		
 The Analysis of Video Conference With ITU Standarization (International Telecommunication Union) That Joining in Inherent At Bandar Lampung University Maria Shusanti F, Happy Reksa		

 26. The E-internal audit iso 9001:2008 based on accreditation form assessment matrix in study program for effectiveness of monitoring accreditation Marzuki, Maria Shusanti F
27. The Developing Of e-Consultations For Effectiveness of Mentoring Academy Ahmad Cucus, Endang K
 The Evaluation of information system performance in higher education case study with EUCS model at bandar lampung university Reni Nursyanti, Erlangga.
 The Analysis Of History Collection System Based On AndroidSmartphone With Qr Code Using Qr CodeCase Study: Museum Lampung Usman Rizal, Wiwin Susanty, Sutrisno
 30. Application of Complaint Handling by Approach Model of ISO 10002 : 2004 to Increase Complaint Services Agus Sukoco and Yuthsi Aprilinda.
 Towards Indonesian Cloud Campus Taqwan Thamrin, Iing Lukman, Dina Ika Wahyuningsih
32. Bridging Router to ADSL Modem for Stability Network Connection Arnes Yuli Vandika and Ruri Koesliandana
 33. The Effect of Use Styrofoam for Flexural Characteristics of Reinforced Concrete Beams Yasser , Herman Parung, M. Wihardi Tjaronge, Rudy Djamaluddin 261
34. The Estimation Of Bioethanol Yield From Some Cassava Variety M.C. Tri Atmodjo
 35. Effect of Superficial Velocity of Pressure Difference on The Separation of Oil And Water by Using The T-Pipe Junctionl Kms. Ridhuan and Indarto
 36. The use of CRM for Customer Management at Cellular Telecommunications Industry Ayu Kartika Puspa
 37. Indonesian Puslit (Centre Of IT Solution) Website Analysis Using Webqual For Measuring Website Quality Maria Shusanti Febrianti and Nurhayati.
 The E-internal audit iso 9001:2008 based on accreditation form assessment matrix in study program for effectiveness of monitoring accreditation Marzuki, Maria Shusanti F

2 nd International Conference on Engineering and Technology Development	ISSN 2301-6590
(ICETD 2013)	
Universitas Bandar Lampung	
Faculty of Engineering and Faculty of Computer Science	

 Enhancing Quality Software Through CMMI-ISO 9001:2008and ISO 9126 Agus Sukoco
 Value Analysis Of Passenger Car Equivalent Motorcycle (Case Study Kartini Road Bandar Lampung) Juniardi, Aflah Efendi
 Alternative Analysis Of Flood Control Downstream Of Way Sekampung River Sugito, Maulana Febramsyah.
 Analysis Of Fitness Facilities And Effective Use Of Crossing Road Juniardi, Edi Haryanto
 Study On Regional Development Work Environment Panjang Port Lands In Support Bandar Lampung City As A Service And Trade Ir. A. Karim Iksan, MT, Yohn Ferry.
44. Analytical And Experimental Study Bamboo Beam ConcreteHery Riyanto, Sugito, Juli
 45. Comparative Analysis Of Load Factor Method Static And Dynamic Method (Case Study Akdp Bus Route Rajabasa - Bakauheni) A. Ikhsan Karim, MT., Ahmad Zulkily
 Optimization Utilization Of Water Resourcesdam Batutegi Using Method Of Linear Program Aprizal,HeryFitriyansyah
 47. Characteristics Generation Traffic Patterns And Movement In Residential Area (Case Study Way Kandis Residential Bandar Lampung) Fery Hendi Jaya, Juniardi,
 Use Study On Slight Beam Reinforced Concrete Floor Platein Lieu Of Scondary Beam Hery Riyanto, Sugito, Lilies Widodjoko, Sjamsu Iskandar
 Observation Of The Effect Of Static Magnetic Field 0.1 Mt On A-Amylase Activity In Legume Germination Rochmah Agustrina, Tundjung T. Handayani, and Sumardi
 50. Effectiveness Analysis Of Applications Netsupport School 10 Based Iso / Iec 9126-4 Metrics Effectiveness Ahmad Cucus, Nelcy Novelia
51. Omparative Performance Analysis Of Banking For Implementing Internet Banking Reza Kurniawan

OBSERVATION OF THE EFFECT OF STATIC MAGNETIC FIELD 0.1 MT ON α-AMYLASE ACTIVITY IN LEGUME GERMINATION

Rochmah Agustrina¹, Tundjung T. Handayani, and Sumardi ¹Biology Departement, Scholl of Mathematics and basic Sciences University of Lampung, Jln. Soemantri Brojonegoro 1, Bandar Lampung 35145 e-mail : rochmah@unila.ac.id

Abstract-*Studies on the influence of magnetic fields on plant growth and development provide a promising prospect in the field of agricultural engineering. But there are still many obstacles encountered, because the plant response to the magnetic field treatment varies depending on the strength, intensity, and duration of exposure. Type, age, as well as parts of the plant are used also gives a different response to magnetic field treatment. Previous studies showed that a magnetic field can increase the germination of some seeds of agricultural crops. In this study, The entire treatment duration of exposure 0.1 mT magnetic field causes an increase in \alpha-amylase activity in all legume seedlings observed. Fluctuations in enzyme activity during the germination of kidney bean seedlings are relatively the same as the black turtle bean, as well as enzyme activity in soybean and mung bean. But between the two groups of seedling legumes (kidney beans and black turtle beans vs. Soybeans and green beans) show a very clear difference fluctuations enzyme activity during germination. Keywords: \alpha-amylase, seedling*

Introduction

Plants are organisms that cannot move so that the growth is strongly influenced by local environmental factors. Most people assume that the factors responsible for affecting the growth and development of temperature. plants are: humidity, available nutrients in soil, and light. However, it has long been observed the magnetic field effect on plant growth and developent because the earth is actually a source of natural magnetic fields. Telford (1990) explains that according to Karl Frederick Gauss, magnetic field of the Earth has a close relationship with the rotation of the earth because the earth's magnetic poles close to the Earth's spin axis

Study the influence of magnetic fields on various types of plants have been done long ago. However, the results vary depending on the treatment of the magnetic field (strength, frequency, duration of exposure) (Criveanu, H.R. dan G. Taralunga. 2006 dan Atak et al., 2003) as well as the plant species, age and plant organ treated. As a result, a lot of things about the mechanism and the role of magnetic field in affecting the cell metabolism in plant tissues then expressing its influence on the plant growth and development cannot be fully understood (Atak et al., 2003.). This fact attracts many researchers to continue to study it, mainly due to some of the results of the previous studies provide promising prospects in the use of magnetic fields to improve the plant growth and production (De Souza et al., 2005 and Esitken et al., 2004).

Treatment of the magnetic field is known to be able to increase the germination percentage of various types of plants (Mausavizadeh et al., 2013; Pourakbar and Hatami, 2012; Farashas et all, 2011; Aladjadjiyan et al., 2003; Novitsky et al., 2001), germination rate (Mausavizadeh et al., 2013; Vashist and Nagarajan, 2010; Criveanu, and Taralunga. 2006 and Podlesny et al., 2005) growth rate of seedling (Mausavizadeh et al., 2013 and Agustrina et al., 2012), enzyme activities including α - amylase (Mausavizadeh et al., 2013; Pourakbar and Hatami, 2012; Reddy et al, 2012; Vashist, A., and S. Nagarajan. 2010; Atak et al., 2007, and Pintilie et al, 2006), chlorophyll content (Alikamanoglu et al., 2007; Atak et al., 2007; Pintilie et al, 2006; and Novitsky et al., 2001), and the size of parenchyma cells, vascular bundle cells as well as stomata size of young plants (Agustrina et al. 2011).

All of the results of the studies above indicate that treatment of the magnetic field is able to improve the quality of germination. One of the important enzymes in the germination process is α -amylase as instrumental in deciphering the food reserves in the cotyledons polysaccharide into simpler compounds. α -amylase activity during germination induced by an increase in water content at the beginning of seed germination (Vashisth and Nagarajan, 2010). The enzyme α -amylase activity in seeds were germinated breaks the glycoside bonds in the middle of the polysaccharide chain. Their activity creates a mixture of glucose, maltose, isomaltose and oligo-saccharides. Betaamylase then tears off the rest of the maltose from the non-reductive end of the polysaccharide chain (Rochalska and Orzeszko-Rywka, 2005). The beginning process of decomposition of food reserves in the seed source is vital, because the availability of carbon compounds is essential for simple germination metabolic processes so that individuals eventually grow new plants.

This paper report the study study done on the observation of α -amylase activity in the 4 types of 3 legume species: kidney bean and turtle black bean (Phaseolus vulgari), soybean (Glycine max), and mung beans (Vigna radiata) treated with 0.1 mT magnetic field with different long exposure long exposure to determine which treatment is most appropriate for improving quality (vigor) seedlings

Materials and Methods

1. Seed Germination

Legume seeds used in this study were kidney bean and black bean (Phaseolus vulgaris), soy bean (Glycine max), and mung bean (Vigna radiata) were collected from several plant breeding institutions. Seeds were selected for uniform size and shape. They were placed in petridish, given a sufficient water before treated by 0.1 mT magnetic field with exposure duration of: 7'48 "; 11'44", and 15'36 ". controls were magnetically As unexposed-seeds. The petridishes then were placed in germinator until the seedlings are ready to be used for measurements of α -amylase enzyme activity.

2. α-amylase activity

Observation α -amylase activity was done on the basis of seedling age (0.5 to 7 days) and length of hypocotyl (1, 1)3, 5, 7, and 9 cm). In order to measure the activity of α -amylase, the following reaction was set up. a) Sample analysis: 250 μ 1 250 μ 1 enzyme and substrate μ 1 (0.1% starch) were incubated in 30 ° C for 10 minutes. The reaction was stoped by adding 250 µl HCl, 250 µ l iodine solution, and 4 ml of distilled water. b) Control analysis: 250 μ l enzyme activity was inactivated by adding 250 µl of 1 N HCl before incubation at 30 ° C for 10 minutes. Into the tube then the substrate (0.1% starch) 25 µ 1, 250 µ 1 iodine solution, and 4 ml of distilled water were added.

The α -amylase enzyme activity wasas then measured at Abs575

Results

Water plays a key role in the germination process. Germination is the process of growth of the zygote which begins with the process of water imbibition. Water status changes in enzyme activity in seeds stimulate germination, increased uptake of oxygen as a sign of an increase in seed respiration and metabolism. Germination process is completed with radicle protrusion through the seed coat and adsorption of water and oxygen rapidly increases (Vashisth and Nagarajan, 2010). Alpha amylase plays an important role in hydrolyzing the endosperm starch into sugars, which provide the energy for the growth of roots and shoots (Kaneko et al., 2002).

Studies on seed germination of various crops showed that magnetic field treatment can accelerate the hydration of seeds and improve seedling fresh weight, germination percentage (Bautista, 2013 and Reddy et al 2013) seedling dry weight (Pourakbar and Hatami, 2012), and the germination rate (De Souza et al., 2006).





Control

7'48"

11'44"

15'36"





Figure 1. Changes in the activity of α amylase in legum seedling of *Phaseolus vulgaris*: (A) kidney bean and (B) black turtle bean, (C) *Glycine max*: white soybean, and (D) *Vigna radiata*: mung bean within7 days of germination







Figure 2. Changes in the activity of α amylase in legum seedling of *Phaseolus vulgaris*: (A) kidney bean and (B) black turtle bean, (C) *Glycine max*: white soybean, and (D) *Vigna radiata*: mung bean measuerd based on length of hypocotyl

Based on several other studies Morejon et al. (2007) explains in his paper that magnetic field exposure to normal water (water+ions) changes some of its physical and chemical properties such as: surface tension, conductivity,

solubility of salts, refractive index and By application of magnetically pH. treated water (MTW) in pine seed germination, he showed that it germination percentage. improves The result leads to a hypothesis that MTW properties as mwntioned above make it much easier to can penetrate inner parts of the seed. Several other theories have also been proposed to explain the mechanism of magnetic field effect in plant germination metabolisma, including biochemical changes or altered enzyme activities (Majd and Shabrangi, 2009). Garcia and Arza (2001) carried out an experiment study on water absorption by lettuce seeds previously treated in a stationary magnetic field of 1 to 10 mT. They reported an increase in water uptake rate due to the applied magnetic field, which may be the explanation for the increase in the germination seed of treated lettuce seeds. It seems changes in intracellular levels of ca2+ and in other ionic current density across cellular membrane (Lyle et al., 1991) cause alteration in osmotic pressure and changes in capacity of cellular tissues to absorb water (Garcia and Arza, 2001)

Increase in water status during seed imbibition in maieze seeds as a result of magnetic field treatment is detected early appearance of hydration water, greater amount of cytoplasmic bulk water. hydration water. molecular mobility of cytoplasmic bulk water, hydration water of macromolecules. This early hydration of macromolecules as well as greater activities of membranes and germination enzymes during were responsible for Quicker germination of seeds Vashisth and Nagarajan (2010). These results are consistent with the results of our study on the treatment of the magnetic field exposure to the amylase activity in germination of legume seeds. All of treatments of duration of exposure of magnetic field seems increased the

activity of α -amylase in all legume seedling observed.

The activity of α -amylase in legume seedlings measured both by seedling age (Fig. 1) and length of hypocotyl (Fig.2) is greater in seedlings from treated seeds with magnetic fields exposure than from untreated seeds or controls. Moreover, in the figure above, it can be seen that the entire treatment of exposure duration of magnetic field in the early germination increased enzyme activity. The fact lead to the hypothesis that the entire treatment duration of exposure of magnetic field in the present study has the same effectiveness in increasing the activity of α -amylase during germination.

The enzyme α -amylase known has a substantial role in the degradation of food reserves of the seedling during germination. In the present study, increase in enzyme activity has been seen since the germination: 0.5 davs early of germination (Fig. 1) or at the time of hypocotyl length only reaches about 1 cm (Fig.2) which is in accordance with the results of other studies on the germination enzyme activities in various magneticallytreated seeds. The increase in activity of germination enzymes activity is then followed with the improvement of seedling performance or vigor such as the early onset of germination, germination percentage, seedling growth rate, as well as fresh and dry weight (Mausavizadeh et al., 2013; Pourakbar and Hatami, 2012; Vashisth and Nagarajan, 2010, and Atak et al., 2007). These results strengthen the opinion Reddy et al (2012) that higher enzyme activity in magnetic field treated seeds could be triggering the fast germination and early vigor of seedlings.

The enzym α-amylase activityduringgerminationfluctuationsin enzyme activityduring

germination appear to be relatively similar in kidney bean seedlings and black turtle bean seedlings, it is also the α -amylase activity between white soybean seedlings and mung bean seedlings. But when we compare the α -amylase enzyme activity among legume seelings of *Phaseolus* sp., *Glycine* sp., and Vigna sp., seems very different. This result proves that plants respond differently to the magnetic field treatment, among other things, depending on the plant species.

On Phaseolus, the increase in enzyme α -amylase activity in seedlings from magnetically-treated seeds of mung bean seems greater than those of from white soybean. This is clearly observed when the enzyme activity measurement in seedling done base on the length of hypocotyl. The comparison of α -amylase activity in seedling from magneticallytreated seeds of Phaseolus sp. to other seedlings from Glycine sp., and Vigna sp., showed that the fluctuation of α -amylase seedlings during germination activity magnetically-treated from seeds of Glycine sp., and Vigna sp. are greater (Fig. 1). The observation result of the present study indicates that the highest increase in α -amylase activity as a result of magnetic field exposure in early seed germination is detected in seedlings from magnetically-treated seeds of mung bean (Phaseolus vulgaris).

Conclusion

The present study suggest that the duration of exposure treatmen of magnetic field 0,1 mT from 7'48 "; 11'44", and 15'36 improve the activity of enzyme α -amylase during germination. The discussion on the pereset study leads to the assumption: first, magnetic field change the chemically process in plant

cell system or the enzymes properties (Morejon et al., 2007 and Lyle et al., 1991) and second, the magnetic field changes the properties of water so it become easier to penetrate and hydtrate the seed cells (Reddy et al., 2012; Vashisth dan Nagarajan, 2010). The magnetic field treatment in early germination of legume seeds increases in the activity of enzym α -amylase due to the high increase in water status of seedlings (Vashisth and Nagarajan, 2010) then is followed by improvement of seedling performance or vigor.

As a result, the application of magnetic field on the seed would not only useful as a methods to improve the seedling vigor which is important in agriculture engineering to enhance crop production but as suggested by Reddy et al., (2012) it may also provides and excelent sources of α -amylase, extensively used industrially.

References

Aladjadjiyan, Anna and Teodora Ylieve. 2003. Influence of stationary magnetic field on the early stages of the development of tobacco seeds (*Nicotiana tabacum* L.). Journal Central Europian Agriculture. Vol. 4. No. 2. p131-137

Alikamanoglu, S., Yoycili., A. Atak, and A. Rzakoulieve. 2007. Effect of magnetic field and gamma radiation on *Paulowinia tomntosa* tissue culture. Biotechnology and Biotechnological equiptment. Vol 21(1): 49-53

Agustrina, R., T. T. Handayani, S. Wahyuningsih, dan O. Prasetya. 2012. Pertumbuhan Tanaman Tomat (*Lycopersicum Esculentum* Mill.) di Bawah Perlakuan Medan Magnet 0,2 mT. In the Proceedings of the 2012 SNSMAIP III, pp: 277-281 Agustrina, R., T. Tripeni, dan E. Ernawiati. 2011. Anatomi Kecambah Tomat Yang Diberi Perlakuan Medan Magnet 0,2 mT. In In the Proceedings of the 2012 Seminar Nasional Sains & Teknologi - IV, pp: 632-646

Atak C., O. Celik, A. Olgun1, S. Alikamanoglu, A. Rzakoulieva. 2007. Effect of Agnetic Field on Peroxidase Activities of Soybean Tissue Culture. BIOTECHNOL. & BIOTECHNOL. EQ. 21 (2): 166-171

Atak, C., O. Emiroglu, S. Alimakonoglu, and A. Rzakoulieva. 2003. Stimulation of regeneration by magnetic field in soybean (*Glycine max* L. Merrill) tissue cultures. Journal Of Cell And Molecular Biology. 2: 113 -119.

Bautista, R.O.D., V.Y. Victoria, E.M. Ragragio, M.J.M. Deluna, and G.L. Sia Su, 2013. Effect ov Varying Magnetic Frequencies to The *Oryza sativa* L. Var. 128 Seed Germination and Seedling Development. Journal of Applied Phytotechnology in Environmental Sanitation. 2(1): 1-7.

Criveanu, H.R. dan G. Taralunga. 2006. Influence of magnetic fields of variable intensity on behaviour of some medicinal plants. Journal of Central European Agricultura. Vol 7 (4): 643-648.

De Souza, A., D. García, L. Sueiro, L. Licea and E. Porras. 2005. Pre-sowing magnetic treatment of tomato Seeds: effects on the growth and yield of plants cultivated late in the season. Spanish Journal of Agricultural Research. 3(1), 113-122

Esitken, A dan M. Turan. 2004. Alternating magnetic field effects on yield and plant nutrient element composition of strawberry (*Fragaria xananassa* cv. Camarosa). Acta Agriculture Scandinavica, B, Vol 54 No 3 p.135-139 Farashah, H. D., R. T. Afshari, F. Sharifzadeh, S. Chavoshinasab. 2011. Germination improvement and α -amylase and β -1,3-glucanase activity in dormant and nondormant seeds of Oregano (*Origanum vulgare*). AJCS 5(4): 421-427

Garcia, F. and L. I., Arze. .2001. Influence of a Stationary Magnetic Field on water relations in lettuce seeds. Part I: Theoretical considerations, Bioelectromagnetics, Vol. 22, No. 8, pp. 589-595.

Kaneko, M. Itoh H, Ueguchi-Tanaka M, Ashikari M, Matsuoka M. 2002. The α amylase induction in Endosperm during Rice Seed Germination Is Caused by Gibberellin Synthesized in Ephithelium. Plant Physiology. 168: 1264-1270.

Lyle DB, Wang X, Ayotte RD, Sheppard AR, Adey WR. 1991. Calcium uptake by leukemic and normal tlymphocytes exposed to low frequency magnetic fields. Bioelectromagnetics **12**, 145-156.

Majd, A.and A. Shabrangi. 2009. Efect of Seed Pretreatment by Magnetic Fields on Seed Germination and Ontogeny Growth of Agricultural Plants. In the Proceeding of 2009 Electromagnetics Research Symposium, Beijing, China, 1137-1141

Mousavizadeh, S.J., S. Sedaghathoor, A. Rahimi, H. Mohammadi. 2013. Germination parameters and peroxidase activity of lettuce seed under stationary magnetic field . Int. J. Biosci Vol. 3, No. 4, p. 199-207

Morejón, L.P., J.C.Castro Palacio, L.Velázquez Abad, and A.P. Govea. 2007. Stimulation of Pinus tropicalis M. seeds by magnetically treated water. Int. Agrophysics. 21, 173-177 2nd International Conference on Engineering and Technology Development (ICETD 2013) Universitas Bandar Lampung Faculty of Engineering and Faculty of Computer Science

Novitsky, Y.I., G.V. Novitskaya, T.K. Kocheshkova., G.A. Nechiporenko., M.V. Dobrovol Skii. 2001. Growth of green onions in a weak permanent magnetic field. Russioan Jurnal of Plant Physiology. 8:709-716

Pintilie, M., L. Oprica, M. Surleac, S. Dragutivan, D.E. Creanga. 2006. Enzyme Activity in Plants Treated with magnetic solution. Rom. Journ. Phys., Vol. 51, Nos. 1–2, P. 239–244

Podlesny, J., L.E. Misiak, A. Podlesna, dan S. Pietruszewski. 2005. Concentration of free radicals in pea seeds after pre-sowing treatment wirth magnetic field. International Agrophysics. Vol 19. No.3. pp 243-249.

Pourakbar, L. And S. Hatami. 2012. Exposure of Satureia hortensis L. Seeds to Magnetic Fields: Effect on GerminationGrowth characteristic, and activity of Some wnzymes. Journal of Stress Physiology & Biochemistry. Vol 8. No. 21: 191-198

Reddy, K.V.' S. Raisha Reshna, S. Jareema, and M. Nagaraju. 2012. Exposure of Greengram Seeds (Vigna radiate var radiata) to static magnetic Field: Effects on germination and α -amylase activity. Re. J. Seed science. 1-9.

Rochalska, M. and Orzeszko-Rywka, A. 2005. Magnetic field treatment improves seed performan**ce**.

Telford, W.M.M, <u>W. M. Telford</u>, <u>R. E.</u> <u>Sheriff</u>. 1990. Applied Geophysics. Sesond edition. Cambridge University Press.,

Vashisth, A. and S. Nagarajan. 2010. Characterization of Water distribution an Activities of EnZymes during Germination in Magnetically-exposed Maize (*Zea mays* L.) Seeds. Indian Journal of Biochemistry and Biophysics. Vol 47. pp. 311-38



9

11 11

-

-

4 4

1-1 1-1

(P) (P) (P)

Ţ

÷

(T) (T)

-

JI. Z.A. Pagar Alam No.26 Labuhan Ratu Bandar Lampung 35142 Phone: +62 721 701463 www.ubl.ac.id Lampung - Indonesia

conveighte02013